



www.support4NICUParents.org

Bibliography on Family-Centered Developmental Care

Altimier L, Phillips R. The Neonatal Integrative Developmental Care Model: Seven Neuroprotective Core Measures for Family-Centered Developmental Care. *Newborn Infant Nurs Rev* 2013; 13(1): 9–22.

American Academy of Pediatrics, Committee on Hospital Care, Institute for Patient- and Family-Centered Care. Patient- and family-centered care and the pediatrician's role. *Pediatrics* 2012; 129(2): 394–404.

American Academy of Pediatrics, Committee on Hospital Care, Institute for Patient- and Family-Centered Care. Patient- and family-centered care and the pediatrician's role. *Pediatrics* 2012; 129(2): 394–404.

American Academy of Pediatrics, March of Dimes, and NANN.; 2014. Baby Steps to Home.

<http://babystepstohome.com/>

Barbosa V. Teamwork in the neonatal intensive care unit. *Phys Occup Ther Pediatr* 2013; 33(1): 5–26.

Barr P. A dyadic analysis of negative emotion personality predisposition effects with psychological distress in neonatal intensive care unit parents. *Psychol Trauma* 2012; 4(4): 347–355.

doi:10.1037/a0024228.

Bernard-Bonnin A, Psychosocial Paediatrics Committee of Canadian Society of Pediatrics, Canadian Paediatric Society. Maternal depression and child development. *Pediatr Child Health* 2004; 9(8): 575–83.

Bialoskurski M, Cox C, Hayes J. The nature of attachment in a neonatal intensive care unit. *J Perinat Neonatal Nurs* 1999; 13(1): 66–77.

Bracht M, O'Leary L, Lee S, O'Brien K. Implementing family-integrated care in the NICU: a parent education and support program. *Adv Neonatal Care* 2013; 13(2): 115–26.

Brett J, Staniszewska S, Newburn M, *et al.* A systematic mapping review of effective interventions for communicating with, supporting and providing information to parents of preterm infants. *BMJ Open* 2011; doi 10.1136/bmjopen-2010-000023.

Browne J. Developmental care for high-risk newborns: Emerging science, clinical application, and continuity from Newborn Intensive Care Unit to community. *Clin Perinatol* 2011; 38: 719–29.

Browne, J, Talmi A. Family-based intervention to enhance infant-parent relationships in the neonatal intensive care unit. *J Pediatr Psychol* 2008; 30(8): 1-11.

Caskey M, Stephens B, Tucker R, Vohr B. Importance of parent talk on the development of preterm infant vocalizations. *Pediatrics* 2011; 128(5): 910–6.

Caskey M, Stephens B, Tucker R, Vohr B. Adult talk in the NICU with preterm infants and developmental outcomes. *Pediatrics* 2014; 133(3):e578–84.

Cho J, Holditch-Davis D, Miles S. Effects of maternal depressive symptoms and infant gender on the interactions between mothers and their medically at-risk infants. *J Obstet Gynecol Neonatal Nurs* 2008; 37(1): 58–70.

Cisneros Moore K, Coker K, DeBuisson A, Swett B, Edwards W. Implementing potentially better practices for improving family-centered care in Neonatal Intensive Care Units: successes and challenges. *Pediatrics* 2003; 111(Supplement E1): e450–60.

Cleveland L. Parenting in the Neonatal Intensive Care Unit. *J Obstet Gynecol Neonatal Nurs* 2008; 37: 666–91.

Conde-Agudelo A, Diaz-Rossello J, Belizan J. Kangaroo mother care to reduce morbidity and mortality in low birth weight infants. *Cochrane Database of Systematic Reviews*. 2003;(2), doi 10.1002/14651858.CD002771.

Cooper L, Gooding J, Gallagher J, Sternesky L, Ledsky R, Berns S. Impact of a family-centered care initiative on NICU care, staff and families. *J Perinatol* 2007; 27(Suppl 2): S32–S37.

Coughlin M. Transformative nursing in the NICU: trauma-informed age-appropriate care. Springer Publishing Company; 2014. 198 p.

Davis R, Davis M, Freed G, Clark S. Fathers' depression related to positive and negative parenting behaviors with 1-year-old children. *Pediatrics* 2011; 27(4): 612–8.

DiScala C, Sege R, Guohua L, Reece R. Child abuse and unintentional injuries- A 10 year retrospective. *Arch Pediatr Adolesc Med* 2000; 154(1): 16–22.

Dunn M, Reilly M, Johnston A, Hoopes R Jr, Abraham M. Development and dissemination of potentially better practices for the provision of family-centered care in neonatology: the family-centered care map. *Pediatrics* 2006; 118: S95–107.

Feldman R, Greenbaum C, Oke S, Golding J. Change in mother-infant interactive behavior: relations to change in the mother, the infant, and the social context. *Infant Behav Dev* 1997; 20(2): 151–63.

Gooding J, Cooper L, Blaine A, Franck L, Howse J, Berns S. Family support and family-centered care in neonatal intensive care unit: Origins, advances, impact. *Semin Perinatol* 2011; 35: 20–8.

Graven S. Early visual development: Implications for the Neonatal Intensive Care Unit and care. *Clin Perinatol* 2011; 38: 671–83.

Gudsnuk K, Champagne F. Epigenetic effects of early developmental experiences. *Clin Perinatol* 2011; 38:703–11.

- Hoffman J. A case of shaken baby syndrome after discharge from the newborn intensive care unit. *Adv Neonatal Care* 2005; 5(3): 135–46.
- Harrison T. Family-centered pediatric nursing care: state of the science. *J Pediatr Nurs* 2010; 25: 335–43.
- Hugill, K, Letherby G, Reid T, Lavender T. Experiences of Fathers Shortly After the Birth of Their Preterm Infants. *Journal of Obstetric, Gynecologic, & Neonatal Nursing* 2013; 42: 655–63.
- Huhtala M, Korja R, Lehtonen L, Haataha L, Lapinleimu P, Rautava P *et al.* Parental psychological well-being and behavioral outcome of very low birth weight infants at 3 years. *Pediatrics* 2012; 129(4): e937–e944.
- Kuo D, Houtrow A, Arango P, Kuhlthau K, Simmons J, Neff J. Family-centered care: current applications and future directions in pediatric health care. *Matern Child Health J* 2012; 16: 297–305.
- Liu WF, Laudert S, Perkins B, Macmillan-York E, Martin S, Graven S. The development of potentially better practices to support the neurodevelopment of infants in the NICU. *J Perinatol* 2007; 27: S48–S74.
- Huhtala M, Korja R, Lehtonen L, Haataha L, Lapinleimu P, Rautava P, *et al.* Parental psychological well-being and behavioral outcome of very low birth weight infants at 3 years. *Pediatrics* 2012; 129(4): e937–44.
- Lehtonen L, Ahlquist-Bjorkroth S, Boukydis Z. Close collaboration with parents training program for implementing family-centered care culture in neonatal intensive care units. Department of Pediatrics at Turku University Hospital; 2013. Available at: <https://files.zotero.net/6844381561/Close%20Collab%20Finland.pdf>.
- Lickliter R. The integrated development of sensory organization. *Clin Perinatol* 2011; 38: 591–603.
- Ludwig S. Poll question: do you know why infants in the Neonatal Intensive Care Unit need neonatal therapy services? *Newborn Infant Nurs Rev* 2013; 13(1): 2–4.
- Maroney D. Recognizing the potential effect of stress and trauma on premature infants in the NICU: How outcomes are affected. *JPerinatol* 2003; 23: 679–83.
- McGrath J, Samra H, Kenner C. Family-centered developmental care practices and research: what will the next century bring? *J Perinat Neonatal Nurs* 2011; 25(2): 165–70.
- Mehler K, Wendrich D, Kissgen R, Roth B, Obertheur A, Pillekamp F, *et al.* Mothers seeing their VLBW infants within 3 h after birth are more likely to establish a secure attachment behavior: evidence of a sensitive period with preterm infants? *J Perinatol* 2011; 31(6): 404–10.
- Melnyk B, Feinstein N, Alpert-Gillis L, Fairbanks E, Crean H, Sinkin R, *et al.* Reducing premature infants' length of stay and improving parents' mental health outcomes with the Creating Opportunities for Parent Empowerment (COPE) neonatal intensive care unit program: a randomized, controlled trial. *Pediatrics* 2006; 118(5): e1414–e1427.
- Milgrom J, Newnham C, Martin P, Anderson P, Doyle L, Hunt R, *et al.* Early communication in preterm infants following intervention in the NICU. *Early Hum Dev* 2013; 89(9): 755–62.

- Milgrom J, Newnham C, Anderson P, Doyle P, Gemmill A, Lee K, *et al.* Early sensitivity training for parents of preterm infants: Impact on the developing brain. *Pediatr Res* 2010; 67(3): 330–5.
- Moon C. The role of early auditory development in attachment and communication. *Clin Perinatol* 2011; 38: 657–69.
- Noble L. Developments in neonatal technology continue to improve infant outcomes. *Pediatr Ann* 2003; 32(9): 595–603.
- O'Brien K, Bracht M, Macdonell K, McBride T, Robson K, O'Leary L, *et al.* A pilot cohort analytic study of Family Integrated Care in a Canadian neonatal intensive care unit. *BMC Pregnancy Childbirth* [Internet]. 2013; 13((Suppl 1)): S12. Available from: <http://www.biomedcentral.com/1471-2494/13/S1/S12>
- Ortenstrand A, Westrup B, Brostrom E, Sarman I, Akerstrom S, Brune T, *et al.* The Stockholm Neonatal Family Centered Care Study: Effects on length of stay and infant morbidity. *Pediatrics* 2010; 125(2): e278–85.
- Petersen M, Cohen J, Parsons V. Family-centered care: do we practice what we preach? *J Obstet Gynecol Neonatal Nurs* 2004; 33(4): 421–7.
- Phillips R. The sacred hour: uninterrupted skin-to-skin contact immediately after birth. *Newborn Infant Nurs Rev* 2013; 13: 67–72.
- Pierrehumbert B, Nicole A, Muller-Nix C, Forcada-Guex M, Ansermet F. Parental post-traumatic reactions after premature birth: implications for sleeping and eating problems in the infant. *Arch Dis Child Fetal Neonatal Ed* 2003; 88(5): 400–4.
- Pitcher J, Schneider L, Drysdale J, *et al.* Motor system development of the preterm and low birthweight infant. *Clin Perinatol* 2011; 38: 605–25.
- Placencia F, McCullough L. Biopsychosocial risks of parental care for high-risk neonates: implications for evidence-based parental counseling. *J Perinatol* 2012; 32(5): 381–6.
- Reynolds L, Duncan M, Smith G. Parental presence and holding in the neonatal intensive care unit and associations with early neurobehavior. *J Perinatol* 2013; 33: 636–41.
- Scher M, Ludington-Hoe S, Kaffashi F, Johnson M, Holditch-Davis D, Loparo K. Neurophysiologic assessment of brain maturation after an eight-week trial of skin-to-skin contact on preterm infants. *Clin Neurophysiol* 2009; 120(10): 1812–8.
- Shonkoff J, Garner A, Siegel B, Dobbins M, Earls M, Garner A, *et al.* The lifelong effects of early childhood adversity and toxic stress. *Pediatrics* 2012; 129(1): e232.
- Smith G, Gutovich J, Smyser C, Pineda R, Newnham C, Tjoeng T, *et al.* Neonatal intensive care unit stress is associated with brain development in preterm infants. *Ann Neurol* 2011; 70(4): 541–9.
- Sturdivant C. A collaborative approach to defining neonatal therapy. *Newborn Infant Nurs Rev* 2013; 13(1): 23–6.

Sullivan R, Perry R, Sloan A, *et al.* Infant bonding and attachment to the caregiver: Insights from basic and clinical science. *Clin Perinatol* 2011; 38: 643–55.

Voos K, Ross G, Ward M, Yohay A, Osorio S, Perlman J. Effects of implementing family-centered rounds (FCR) in a neonatal intensive care unit (NICU). *J Matern Fetal Neonatal Med* 2012; 24(11): 1403–6.

Warre R, O'Brien K, Lee S. Parents as the primary caregivers for their infant in the NICU: benefits and challenges. *NeoReviews* 2014; 15(11): e472–7.

Weis J, Zoffman V, Egerod I. Enhancing person-centred communication in NICU: a comparative thematic analysis. *Nurs Crit Care* 2013 Nov 14; epub ahead of print, doi 10.1111/nicc.12062.

Westrup B, Sizun J, Lagercrantz H. Family-centered developmental supportive care: a holistic and humane approach to reduce stress and pain in neonates. *J Perinatol* 2007; 27(S1): S12–8.

White-Traut R. Reduced healthcare utilization at 6-week corrected age among premature infants after the H-HOPE Mother–Infant Developmental Intervention. *J Obstet Gynecol Neonatal Nurs* 2013; 42(S1): S87.

White-Traut R, Rankin K, Yoder J, Liu L, Vasa R, Geraldo V, *et al.* Influence of H-HOPE intervention for premature infants on growth, feeding progression and length of stay during initial hospitalization. *J Perinatol* 2015; epub. doi: 10.1038/jp.2015.11.